arXiv:1601.04859v1 [cs.IT] 19 Jan 2016

# The Structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes *

N. Annamalai
Research Scholar
Department of Mathematics
Bharathidasan University
Tiruchirappalli-620 024, Tamil Nadu, India
Email: algebra.annamalai@gmail.com

C. Durairajan
Assistant Professor
Department of Mathematics
School of Mathematical Sciences
Bharathidasan University
Tiruchirappalli-620024, Tamil Nadu, India
Email: cdurai66@rediffmail.com

**Proposed running head:** The Structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes

*The first author would like to thank the Department of Science and Technology (DST), New Delhi, India for their financial support in the form of INSPIRE Fellowship (DST Award Letter No.IF130493/DST/INSPIRE Fellowship / 2013/(362) Dated: 26.07.2013) to carry out this work.

**Abstract**

In this paper, we study the algebraic structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes which are $\mathbb{Z}_2[u,v]$-submodules where $u^2 = v^2 = 0$ and $uv = vu$. In particular, we determine a Gray map from $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$ to $\mathbb{Z}_2^{2\alpha+8\beta}$ and study generator and parity check matrices for these codes. Further we study the structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive cyclic codes and constacyclic codes.

*Keywords:* Additive codes; Generator Matrix; Parity-check Matrix.
*2000 Mathematical Subject Classification:* Primary: 94B25, Secondary: 11H31

Corresponding author:

Dr. C. Durairajan
Assistant Professor
Department of Mathematics
Bharathidasan University
Tiruchirappalli-620024, Tamil Nadu, India
E-mail: cdurai66@rediffmail.com

# 1 Introduction

Let $\mathbb{Z}_2$ be the ring of integers modulo 2. Let $\mathbb{Z}_2^n$ denote the set of all binary vectors of lenght $n$. Any non-empty subset of $\mathbb{Z}_2^n$ is a binary code of length $n$ and a subgroup of $\mathbb{Z}_2^n$ is called a binary linear code. In this paper, we introduce a subfamily of binary linear codes, called $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-linear codes.

Additive codes were first defined by Delsarte in 1973 in terms of association schemes [5], [9]. In a translation association scheme, an additive code is generally defined as a subgroup of the underlying abelian group.

Codes over finite rings were studied in [6], [8] and with the remarkable paper by Hammons *et.al*[5] have been studied intensively for the last four decades. A study on mixed alphabet codes has been introduced and some bounds have been presented by Brouwer *et.al.*[1].

Later, $\mathbb{Z}_2\mathbb{Z}_4$-additive codes were generalized to $\mathbb{Z}_2\mathbb{Z}_{2^s}$-additive codes by Aydogu and Siap [2]. For $s \geq 2$, these generalisations to $\mathbb{Z}_2\mathbb{Z}_{2^s}$-additive codes are interesting since they provided good binary codes via Gray maps with rich algebraic structure. Furthermore, the structure of $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$-additive codes and their duals has been determined [3]. In this correspondance, begin inspired by these additive codes, a generalization towards another direction that have a good algebraic structure and provide good binary codes is presented.

The aim of this paper is the study of the algebraic structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes and their dual codes. It is organized as follows. In Section II, we recall the linear codes over the ring $\mathbb{Z}_2 + u\mathbb{Z}_2$ and the linear codes over the ring $\mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2 + uv\mathbb{Z}_2$. In Section III, we study the concept of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes and duality, we determine the generator matrices of the these codes. In section IV, we study the structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive cyclic codes. In section V, we study the structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive constacyclic codes.

# 2 Linear Codes over Chain Rings

Let $R$ be a ring, then $R^n$ is an $R$-module. A submodule of $R^n$ is called $R$-linear code. We denote the Hamming weight of a word as $w_H$ and Lee weight as $w_L$.

The set $\mathbb{Z}_2 + u\mathbb{Z}_2 = \{0, 1, u, 1+u\}$, where $u^2 = 0$, is a commutative finite chain ring of four elements. We denote this ring as $R_1$.

In [7], a $R_1$-linear code $C$ is permutation equivalent to a code with generator matrix

$$G = \begin{bmatrix} I_{l_0} & A & B_1 + uB_2 \\ 0 & uI_{l_1} & uD \end{bmatrix},$$

where $A, B_1, B_2$ and $D$ are matrices over $\mathbb{Z}_2$.

Define $w_L(a+ub) = w_H(b, a \oplus b)$ for all $a, b \in \mathbb{Z}_2$ where the Hamming weight of a word is the number of non-zero coordinates in it.

The definition of the weight immediately leads to a Gray map from $R_1$ to $\mathbb{Z}_2^2$ which can naturally be extended to $R_1^n$:

$\chi_L : (R_1^n, \text{Lee weight}) \to (\mathbb{Z}_2^{2n}, \text{Hamming weight})$ is defined as:

$$\chi_L(\bar{a} + u\bar{b}) = (\bar{b}, \bar{a} \oplus \bar{b}),$$

where $\bar{a}, \bar{b} \in \mathbb{Z}_2^n$ and $\oplus$ denotes addition in $\mathbb{Z}_2$.

Let $R_2$ be the ring $\mathbb{Z}_2[u,v] := \mathbb{Z}_2 + u\mathbb{Z}_2 + v\mathbb{Z}_2 + uv\mathbb{Z}_2$ with $u^2 = 0$, $v^2 = 0$ and $uv = vu$ where $\mathbb{Z}_2 = \{0, 1\}$. Then $R_2$ is a commutative finite chain ring of 16 elements and $a + ub + vc + uvd \in R_2$ is a unit iff $a \neq 0$.

The group of units of $R_2$ is given by $R_2^* = \{1, 1+u, 1+v, 1+uv, 1+u+v, 1+u+v+uv, , 1+u+uv, 1+v+uv\}$. It is also a local ring with the unique maximal ideal $uvR_2$.

A linear code $C$ of lenght $n$ over the ring $R_2$ is an $R_2$-submodule of $R_2^n$.

A non-zero $R_2$-linear code $C$ has a generator matrix which after a suitable permutation of the coordinates can be written in the form:

$$G = \begin{bmatrix} I_{k_0} & A_{01} & A_{02} & A_{03} & A_{04} \\ 0 & uI_{k_1} & uA_{12} & uA_{13} & uA_{14} \\ 0 & 0 & vI_{k_2} & vA_{23} & vA_{24} \\ 0 & 0 & 0 & uvI_{k_3} & uvA_{34} \end{bmatrix} \tag{2.1}$$

where $A_{ij}$ are all matrices over $\mathbb{Z}_2$.

To define the Lee weights and Gray maps for codes over $R_2$. Define $w_L(a+ub+vc+uvd) = w_H(d, a \oplus d, b \oplus d, a \oplus b \oplus d, c \oplus d, a \oplus c \oplus d, b \oplus c \oplus d, a \oplus b \oplus c \oplus d)$ for all $a, b, c, d \in \mathbb{Z}_2$. The definition of the weight immediately leads to a Gray map from $R_2$ to $\mathbb{Z}_2^8$ which can naturally be extended to $R^n$:

$\phi_L : (R_2^n, \text{Lee weight}) \to (\mathbb{Z}_2^{8n}, \text{Hamming weight})$

is defined as:

$$\begin{aligned} \phi_L(\bar{a} + u\bar{b} + v\bar{c} + uv\bar{d}) = (&\bar{d}, \bar{a} \oplus \bar{d}, \bar{b} \oplus \bar{d}, \bar{a} \oplus \bar{b} \oplus \bar{d}, \bar{c} \oplus \bar{d}, \\ &\bar{a} \oplus \bar{c} \oplus \bar{d}, \bar{b} \oplus \bar{c} \oplus \bar{d}, \bar{a} \oplus \bar{b} \oplus \bar{c} \oplus \bar{d}) \end{aligned}$$

where $\bar{a}, \bar{b}, \bar{c}, \bar{d} \in \mathbb{Z}_2^n$.

# 3 $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes

We know that the ring $\mathbb{Z}_2[u]$ is a subring of the ring $R_2$. Being inspired by the structure of $\mathbb{Z}_2\mathbb{Z}_2[u]$-additive codes, we define the following set:

$$\mathbb{Z}_2[u]\mathbb{Z}_2[u,v] = \{(a,b) \mid a \in \mathbb{Z}_2[u] \text{ and } b \in \mathbb{Z}_2[u,v]\}.$$

The set $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$ cannot be endowed with algebraic structure directly. It is not well defined with respect to the usual scalar multiplication by $v \in R_2$. Therefore, this set is not an $R_2$-module. To make it well defined and enrich with an algebraic structure we introduce a new scalar multiplication as follows: We define a mapping

$$\begin{gathered}\eta : R_2 \to \mathbb{Z}_2[u], \\ \eta(a + ub + vc + uvd) = a + ub \\ \text{i.e., } \eta(x) = x \mod v.\end{gathered}$$

It is easy to conclude that $\eta$ is a ring homomorphism. Using this map, we define a scalar multiplication. For $v = (a_0, a_1, \cdots, a_{\alpha-1}, b_0, b_1, \cdots, b_{\beta-1}) \in \mathbb{Z}_2[u]^\alpha \times R_2^\beta$ and $l \in R_2$, we have

$$lv = (\eta(l)a_0, \eta(l)a_1, \cdots, \eta(l)a_{\alpha-1}, lb_0, lb_1, \cdots, lb_{\beta-1}). \tag{3.1}$$

A non-empty subset $C$ of $R_1^\alpha \times R_2^\beta$ is called $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$**-additive code** if it is a $R_2$-submodule of $R_1^\alpha \times R_2^\beta$ with respect to the scalar multiplication defined in Eq.3.1. Then the binary image $\Phi(C) = D$ is called $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-linear code of length $n = 2\alpha + 8\beta$ where $\Phi$ is a map from $R_1^\alpha \times R_2^\beta$ to $\mathbb{Z}_2^n$ defined as

$$\begin{aligned}\Phi(x,y) = (&q_0, q_1, \cdots, q_{\alpha-1}, q_0 + r_0, \cdots, q_{\alpha-1} + r_{\alpha-1}, \\ &d_0, d_1, \cdots, d_{\beta-1}, a_0 + d_0, a_1 + d_1, \cdots, a_{\beta-1} + d_{\beta-1}, \\ &b_0 + d_0, b_1 + d_1, \cdots, b_{\beta-1} + d_{\beta-1}, \\ &a_0 + b_0 + d_0, a_1 + b_1 + d_1, \cdots, a_{\beta-1} + b_{\beta-1} + d_{\beta-1}, \\ &c_0 + d_0, c_1 + d_1, \cdots, c_{\beta-1} + d_{\beta-1}, \\ &a_0 + c_0 + d_0, a_1 + c_1 + d_1, \cdots, a_{\beta-1} + c_{\beta-1} + d_{\beta-1}, \\ &b_0 + c_0 + d_0, b_1 + c_1 + d_1, \cdots, b_{\beta-1} + c_{\beta-1} + d_{\beta-1}, \\ &a_0 + b_0 + c_0 + d_0, \cdots, a_{\beta-1} + b_{\beta-1} + c_{\beta-1} + d_{\beta-1}),\end{aligned}$$

for all $x = (x_0, x_1, \cdots, x_{\alpha-1}) \in R_1^\alpha$ where $x_i = r_i + uq_i$ for $0 \le i \le \alpha - 1$ and $y = (y_0, y_1, \cdots, y_{\beta-1}) \in R_2^\beta$, where $y_j = a_j + ub_j + vc_j + uvd_j$ for $0 \le j \le \beta - 1$.

If $C \subseteq R_1^\alpha \times R_2^\beta$ is a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive code, group isomorphic to $\mathbb{Z}_2^{2l_0} \times \mathbb{Z}_2^{l_1} \times \mathbb{Z}_2^{4k_0} \times \mathbb{Z}_2^{3k_1} \times \mathbb{Z}_2^{2k_2} \times \mathbb{Z}_2^{k_3}$, then $C$ is called a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$**-additive code of type** $(\alpha, \beta; l_0, l_1; k_0, k_1, k_2, k_3)$ where $l_0, l_1, k_0, k_1, k_2$, and $k_3$ are defined above.

## 3.1 Generator matrices of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes

**Theorem 3.1.** *Let $C$ be a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive code of type $(\alpha,\beta;l_0,l_1;k_0,k_1,k_2,k_3)$. Then $C$ is permutation equivalent to a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive code with standard form matrix.*

$$G=\begin{bmatrix} B & T \\ S & A \end{bmatrix} \tag{3.2}$$

*where*

$$B=\begin{bmatrix} I_{l_0} & B_{01} & B_{02} \\ 0 & uI_{l_1} & uB_{12} \end{bmatrix}$$

$$T=\begin{bmatrix} 0 & 0 & 0 & vT_{01} & vT_{02} \\ 0 & 0 & 0 & 0 & uvT_{12} \end{bmatrix}$$

$$S=\begin{bmatrix} 0 & S_{01} & S_{02} \\ 0 & 0 & uS_{12} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$A=\begin{bmatrix} I_{k_0} & A_{01} & A_{02} & A_{03} & A_{04} \\ 0 & uI_{k_1} & uA_{12} & uA_{13} & uA_{14} \\ 0 & 0 & vI_{k_2} & vA_{23} & vA_{24} \\ 0 & 0 & 0 & uvI_{k_3} & uvA_{34} \end{bmatrix}$$

.

*Here, $B_{ij}$ are matrices over $R_1$ and $T_{ij}$ are matrices over $R_2$ for $0\leq i<2$ and $0<j\leq 2$. Further, for $0\leq d<2,\ 0\leq t<4,\ 0<q\leq 4$, $S_{dj}$ and $A_{tq}$ are matrices over $R_2$. Also, $I_{l_w}$ and $I_{k_y}$ are the identity matrices of size $l_w$ and $k_y$, where $0\leq w\leq 1$ and $0\leq y\leq 3$.*

*Proof.* First, by focusing on the first $\alpha$ coordinates which is a projection of the $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive code to a linear $\mathbb{Z}_2[u]$-code and applying the necessary row operations together with column operations(if neessary) we obtain the matrix $B$ as in the upper left corner of the matrixEq.3.2. Next, by focusing on the last $\beta$ coordinates and protecting the form of the matrix $B$, one consider linear codes over the ring $\mathbb{Z}_2[u,v]$ for which the generator matrices are well known to be of the form $A$ as in Eq.3.2. Finally, by protecting both the form of $G$ as in Eq.3.2 which is referred to as the standard form matrix.

□

**Example 3.2.** *Let $C$ be a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive code of type $(2,3;1,1;1,1,1,0)$ with*

*standard generator matrix of $C$ is*

$$\left[\begin{array}{ccc|ccccc} 1 & 1 & 0 & 0 & 0 & 0 & v & v \\ 0 & u & u & 0 & 0 & 0 & 0 & uv \\ \hline 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & u & 0 & u & u & u & u \\ 0 & 0 & 0 & 0 & 0 & v & v & v \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & uv \end{array}\right] \tag{3.3}$$

*and $C$ has $|C| = 2^2 2^1 2^4 2^3 2^2 = 2^{12} = 4096$ codewords.*

## 3.2 Duality of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes

An inner product for two elements $x, y \in R_1^\alpha \times R_2^\beta$ is defined as

$$\langle x, y\rangle = uv\left(\sum_{i=1}^{\alpha} x_i y_i\right) + \sum_{j=\alpha+1}^{\alpha+\beta} x_j y_j \in R_2.$$

Let $D$ be a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$- linear code of length $n$ of type $(\alpha, \beta; l_0, l_1; k_0, k_1, k_2, k_3)$. We denote by $C$ the corresponding additive code, i.e., $C = \Phi^{-1}(D)$. The additive dual code of $C$, denoted by $C^\perp$, is defined in the standard way

$$C^\perp = \{y \in R_1^\alpha \times R_2^\beta \mid \langle x, y\rangle = 0 \text{ for all } x \in C\}$$

The corresponding binary code $\Phi(C^\perp)$ is denoted by $C_\perp$ and is called the $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-dual code of $C$.

The additive dual code $C^\perp$ also an additive code, that is a subgroup of $R_1^\alpha \times R_2^\beta$.

**Theorem 3.3.** *Let $C$ be a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-linear code of type $(\alpha, \beta; l_0, l_1; k_0, k_1, k_2, k_3)$. The $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-dual code $C_\perp$ is then of type $(\alpha, \beta; l_0', l_1'; k_0', k_1', k_2', k_3')$, where*

$$\begin{aligned} l_0' &= \alpha - l_0 - l_1 \\ l_1' &= l_1 \\ k_0' &= \beta - k_0 \\ k_1' &= \beta - k_1 - k_2 \\ k_2' &= k_2 \\ k_3' &= \beta - k_2 - k_3 \end{aligned}$$

## 3.3 Parity-check matrices of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes

As in the classical case, the generator matrix of the dual code is important. In the following theorem, the standard form of the generator matrix of the dual code is presented.

**Theorem 3.4.** *Let $C$ be a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive code of type $(\alpha,\beta;l_0,l_1;k_0,k_1,k_2,k_3)$ with the standard form matrix defined in equation (3.2). Then the generator matrix for the additive dual code $C^\perp$ is given by*

$$H = \begin{bmatrix} \bar{B} & U \\ V & \bar{A}+E \end{bmatrix} \tag{3.4}$$

*where*

$$\bar{B} = \begin{bmatrix} -B_{02}^t + B_{12}^t B_{01}^t & -B_{12}^t & I_{\alpha-l_0-l_1} \\ -uB_{01}^t & uI_{l_1} & 0 \end{bmatrix}$$

$$V = \begin{bmatrix} -T_{02}^t + T_{12}^t B_{01}^t + A_{34}^t T_{01}^t & -T_{12}^t & 0 \\ -uT_{01}^t & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$U = \begin{bmatrix} -uS_{12}^t A_{01}^t + u(B_{12}^t S_{01}^t - S_{02}^t) & -uS_{12}^t & 0 & 0 & 0 \\ -uvS_{01}^t & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$E = \begin{bmatrix} T_{12}^t S_{01} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\bar{A} = \begin{bmatrix} l & -um & -uA_{12}^t & -uA_{23}^t & uI_{\beta-k_1-k_2} & 0 \\ -vp & -vA_{12}^t & vI_{k_2} & 0 & 0 & \\ -uvA_{01}^t & uvI_{\beta-k_2-k_3} & 0 & 0 & 0 & \end{bmatrix}$$

*where*

$l = -A_{04}^t + A_{01}^t(A_{14}^t + A_{13}^t A_{34}^t + A_{24}^t A_{12}^t + A_{12}^t A_{23}^t A_{34}^t) + A_{02}^t(A_{24}^t + A_{23}^t A_{34}^t) + A_{03}^t A_{34}^t$

$k = A_{14}^t + A_{13}^t A_{34}^t + A_{24}^t A_{12}^t + A_{12}^t A_{23}^t A_{34}^t$

$n = A_{24}^t + A_{23}^t A_{34}^t$

$m = A_{03}^t + A_{02}^t A_{23}^t + A_{01}^t A_{12}^t$

$p = A_{02}^t + A_{01}^t A_{12}^t$

*Proof.* It is easy to check that $HG^t = 0$. Besides, since we have $|C||C^\perp| = 2^{2\alpha+8\beta}$ by checking the type of the matrix $H$ we conclude that the rows of $H$ not only are orthogonal to the rows of $G$ but also generate the all code $C^\perp$. Hence, $H$ is the desired matrix. □

**Example 3.5.** *Let $C$ be a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$- additive code of type $(2,3;1,1;1,1,1,0)$ with the generator matrix given in Equation (3.3). Then, we can write the parity-check matrix of $C$ as follows:*

$$H = \left[\begin{array}{ccc|ccccc} 1 & 1 & 1 & u & u & 0 & 0 & 0 \\ u & u & 0 & uv & 0 & 0 & 0 & 0 \\ \hline 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ u & 0 & 0 & u & u & u & u & 0 \\ 0 & 0 & 0 & v & v & v & 0 & 0 \\ 0 & 0 & 0 & 0 & uv & 0 & 0 & 0 \end{array}\right].$$

*It is clear that is of type $(2,3;0,1;2,1,1,2)$ and has $|C^{\perp}| = 2^1 2^8 2^3 2^2 2^2 = 2^{16} = 65536$ codewords.*

# 4 The Structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive cyclic code

In this section, we introduce the definition of a additive cyclic code and some algebraic structure.

Let $R_{\alpha,\beta}[x] = \dfrac{R_1[x]}{< x^{\alpha}-1 >} \times \dfrac{R_2[x]}{< x^{\beta}-1 >}$.

A code $C$ is cyclic if and only if its polynomial representation is an ideal.

A additive code $C$ is called a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$**-additive cyclic code** if any cyclic shift of a codeword is also a codeword.

i.e., $(a_0, a_1, \cdots, a_{\alpha-1}, b_0, b_1, \cdots, b_{\beta-1}) \in C$

$\Rightarrow (a_{\alpha-1}, a_0, \cdots, a_{\alpha-2}, b_{\beta-1}, b_0, \cdots, b_{\beta-2}) \in C$.

**Theorem 4.1.** *If $C$ is any $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive cyclic code, then $C^{\perp}$ is also cyclic.*

*Proof.* Let $C$ be any $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive cyclic code.

Suppose $x = (a_0, a_1, \cdots, a_{\alpha-1}, b_0, b_1, \cdots, b_{\beta-1}) \in C^{\perp}$,

for any codeword

$y = (d_0, d_1, \cdots, d_{\alpha-1}, e_0, e_1, \cdots, e_{\beta-1}) \in C$

we have

$\langle x, y\rangle = uv\left(\sum_{i=0}^{\alpha-1} a_i d_i\right) + \sum_{j=0}^{\beta-1} b_j e_j = 0.$

Let $S$ is a cyclic shift, and $j = lcm(\alpha, \beta)$. Then we have

$S(x) = (a_{\alpha-1}, a_0, \cdots, a_{\alpha-2}, b_{\beta-1}, b_0, \cdots, b_{\beta-2})$ and $S^j(y) = y$ for any $y \in C$.

Since $C$ be any $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive cyclic code, So we have

$S^{j-1}(y) = (d_1, d_2, \cdots, d_{\alpha-1}, d_0, e_1, e_2, \cdots, e_{\beta-1}, e_0) \in C$.

Hence

$$\begin{aligned}
0 &= \langle x, S^{j-1}(y)\rangle \\
&= uv(a_0d_1 + a_1d_2 + \cdots + a_{\alpha-2}d_{\alpha-1} + a_{\alpha-1}d_0) \\
&+ (b_0e_1 + b_1e_2 + \cdots b_{\beta-2}e_{\beta-1} + b_{\beta-1}e_0) \\
&= uv(a_{\alpha-1}d_0 + a_0d_1 + a_1d_2 + \cdots + a_{\alpha-2}d_{\alpha-1}) \\
&+ (b_{\beta-1}e_0 + b_0e_1 + b_1e_2 + \cdots b_{\beta-2}e_{\beta-1}) \\
&= \langle S(x), y\rangle
\end{aligned}$$

Therefore, we have $S(x) \in C^{\perp}$, so $C^{\perp}$ is a cyclic code. □

Let $C$ be a $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive cyclic code, for any codeword

$c = (a_0, a_1, \cdots, a_{\alpha-1}, b_0, b_1, \cdots, b_{\beta-1}) \in C$ can be representation with a polynomial, that is,

$c(x) = (a(x), b(x)) \in R_{\alpha,\beta}[x]$

Similarly, we introduce a new scalar multiplication. Now, we have the following scalar multiplication:

for $c_1(x) = (a_1(x), b_1(x)) \in R_{\alpha,\beta}[x]$ and $h(x) = p(x)+uq(x)+vr(x)+uvs(x) \in R_2[x]$, define

$h(x)c_1(x) = ((p(x) + uq(x))a_1(x), h(x)b_1(x))$.

Now, we define the homomorphism mapping:

$$\Psi : R_{\alpha,\beta}[x] \to R_2[x]$$

$$\Psi(c(x)) = \Psi(a(x), b(x)) = b(x).$$

It is clear that Image($\Psi$) is an ideal in the ring $\dfrac{R_2[x]}{< x^{\beta} - 1 >}$ and ker($\Psi$) is also an ideal in $\dfrac{R_1[x]}{< x^{\alpha} - 1 >}$. And note that

$Image(\Psi) = \{b(x) \in R_2[x] : (a(x), b(x)) \in R_{\alpha,\beta}[x]\}$

$ker(\Psi) = \left\{(a(x), 0) \in R_{\alpha,\beta}[x] : a(x) \in \dfrac{R_1[x]}{< x^{\alpha} - 1 >}\right\}$.

We have

$Image(\Psi) =< g(x) + up_1(x) + vp_2(x) + uvp_3(x), ua_1(x) + vq_1(x) + uvq_2(x), va_2(x) + uvr_1(x), uva_3(x) >$

where $g(x), p_1(x), p_2(x), p_3(x), a_1(x), q_1(x), q_2(x), a_2(x), r_1(x), a_3(x) \in \dfrac{R_2[x]}{< x^{\beta} - 1 >}$ and

$a_3(x) \mid a_2(x) \mid a_1(x) \mid g(x) \mid x^{\beta} - 1 \ mod\ 2$,

$a_1(x) \mid p_1(x) \left(\dfrac{x^{\beta} - 1}{g(x)}\right)$,

$a_2(x) \,|\, q_1(x) \left(\frac{x^\beta - 1}{a_1(x)}\right)$,
$a_2(x) \,|\, p_2(x) \left(\frac{x^\beta - 1}{g(x)}\right) \left(\frac{x^\beta - 1}{a_1(x)}\right)$,
$a_3(x) \,|\, r_1(x) \left(\frac{x^\beta - 1}{a_2(x)}\right)$,
$a_3(x) \,|\, q_2(x) \left(\frac{x^\beta - 1}{q_1(x)}\right) \left(\frac{x^\beta - 1}{a_1(x)}\right)$.
$a_3(x) \,|\, p_3(x) \left(\frac{x^\beta - 1}{g(x)}\right) \left(\frac{x^\beta - 1}{a_2(x)}\right) \left(\frac{x^\beta - 1}{a_1(x)}\right)$.

Similarly,

$$ker(\Psi) =< ((f(x) + ul(x), ub(x)), 0) >$$

where $f(x), l(x), b(x) \in \frac{R_1[x]}{< x^\alpha - 1 >}$ and
$b(x) \,|\, f(x) \,|\, x^\alpha - 1$, $b(x) \,|\, l(x) \left(\frac{x^\alpha - 1}{f(x)}\right)$.

According to the fundamendal theorem of homomorphism, we have
$\frac{C}{ker(\Psi)} \cong< g(x) + up_1(x) + vp_2(x) + uvp_3(x), ua_1(x) + vq_1(x) + uvq_2(x), va_2(x) + uvr_1(x), uva_3(x) >$.

Hence, we have $((h(x) + ut(x), uc(x)), (g(x) + up_1(x) + vp_2(x) + uvp_3(x), ua_1(x) + vq_1(x) + uvq_2(x), va_2(x) + uvr_1(x), uva_3(x))) \in C$,

where $\Psi((h(x) + ut(x), uc(x)), (g(x) + up_1(x) + vp_2(x) + uvp_3(x), ua_1(x) + vq_1(x) + uvq_2(x), va_2(x)+uvr_1(x), uva_3(x))) = (g(x)+up_1(x)+vp_2(x)+uvp_3(x), ua_1(x)+vq_1(x)+uvq_2(x), va_2(x) + uvr_1(x), uva_3(x))$.

By these discussion, it is easy to see that any $\mathbb{Z}_2[u]\mathbb{Z}_2[u, v]$-additive cyclic code can be generated by two elements of the form $((h(x) + ut(x), uc(x)), (g(x) + up_1(x) + vp_2(x) + uvp_3(x), ua_1(x) + vq_1(x) + uvq_2(x), va_2(x) + uvr_1(x), uva_3(x)))$ and $((f(x) + ul(x), ub(x)), 0)$.

**Corollary 4.2.** *Let $C$ be a $\mathbb{Z}_2[u]\mathbb{Z}_2[u, v]$-additive cyclic code. Then $C$ is an ideal in $R_{\alpha,\beta}[x]$ which can be generated by*

$C = (((f(x) + ul(x), ub(x)), 0), ((h(x) + ut(x), uc(x)), (g(x) + up_1(x) + vp_2(x) + uvp_3(x), ua_1(x) + vq_1(x) + uvq_2(x), va_2(x) + uvr_1(x), uva_3(x))))$,

*where* $a_3(x) \,|\, a_2(x) \,|\, a_1(x) \,|\, g(x) \,|\, x^\beta - 1$ *mod* 2,
$a_1(x) \,|\, p_1(x) \left(\frac{x^\beta - 1}{g(x)}\right)$,
$a_2(x) \,|\, q_1(x) \left(\frac{x^\beta - 1}{a_1(x)}\right)$,
$a_2(x) \,|\, p_2(x) \left(\frac{x^\beta - 1}{g(x)}\right) \left(\frac{x^\beta - 1}{a_1(x)}\right)$,

$a_3(x) \mid r_1(x)\left(\frac{x^\beta-1}{a_2(x)}\right)$,
$a_3(x) \mid q_2(x)\left(\frac{x^\beta-1}{q_1(x)}\right)\left(\frac{x^\beta-1}{a_1(x)}\right)$.
$a_3(x) \mid p_3(x)\left(\frac{x^\beta-1}{g(x)}\right)\left(\frac{x^\beta-1}{a_2(x)}\right)\left(\frac{x^\beta-1}{a_1(x)}\right)$,
$b(x) \mid f(x) | x^\alpha-1$, $b(x) \mid l(x)\left(\frac{x^\alpha-1}{f(x)}\right)$,
$c(x) \mid h(x) \mid x^\alpha-1$ *and* $c(x) \mid t(x)\left(\frac{x^\alpha-1}{h(x)}\right)$.

# 5 $(1+u, 1+uv)$-additive constacyclic codes over $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$

In this section, we introduce the definition of a additive constacyclic code and some algebraic structure.

For fixed units $1+u \in R_1$ and $1+uv \in R_2$. The shift operator $\tau$ on $R_1^\alpha \times R_2^\beta$ is defined to be

$$\tau(a_0, a_1, \cdots, a_{\alpha-1}, b_0, b_1, \cdots, b_{\beta-1}) = ((1+u)a_{\alpha-1}, a_0, \cdots, a_{\alpha-2}, (1+uv)b_{\beta-1}, b_0, \cdots, b_{\beta-2}).$$

A $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive code $C$ is said to be $(1+u, 1+uv)$**-additive constacyclic code** if the code is inveriant under the shift $\tau$, that is $\tau(C) = C$.

Let $\sigma$ be the cyclic shift. For any positive integer $s$, let $\sigma_s$ be the quasi-shift given by

$\sigma_s(a^{(1)}|a^{(2)}|\cdots|a^{(s)}) = \sigma(a^{(1)})|\sigma(a^{(2)})|\cdots|\sigma(a^{(s)})$,

where$a^{(1)}, a^{(2)}, \cdots, a^{(s)} \in \mathbb{Z}_2^n$ and $|$ denotes the usual vector concatenation. A binary quasi-cyclic code $D$ of index $s$ and lenght $n$ is a subset of $\mathbb{Z}_2^n$ such that $\sigma_s(D) = D$.

**Lemma 5.1.** *Let $\Psi$ be a Gray map defined as above and let $\tau$ be the $(1+u, 1+uv)$-constacyclic shift on $R_1^\alpha \times R_2^\beta$. Then $\Psi\tau = \sigma_2\Psi$.*

*Proof.* Let $w = (x, y) = (x_0, x_1, \cdots, x_{\alpha-1}, y_0, y_1, \cdots, y_{\beta-1}) \in R_1^\alpha \times R_2^\beta$.

Let $x_i = r_i + uq_i$ and $y_j = a_j + ub_j + vc_j + uvd_j$ where $x = (x_0, x_1, \cdots, x_{\alpha-1})$ and $y = (y_0, y_1, \cdots, y_{\beta-1})$ for $0 \le i \le \alpha-1,\ 0 \le j \le \beta-1$. From definitions, we have

$$\begin{aligned}\Phi(w) = (&q_0, q_1, \cdots, q_{\alpha-1}, q_0 + r_0, q_1 + r_1, \cdots, q_{\alpha-1} + r_{\alpha-1},\\ &d_0, d_1, \cdots, d_{\beta-1}, a_0 + d_0, a_1 + d_1, \cdots, a_{\beta-1} + d_{\beta-1},\\ &b_0 + d_0, b_1 + d_1, \cdots, b_{\beta-1} + d_{\beta-1},\\ &a_0 + b_0 + d_0, a_1 + b_1 + d_1, \cdots, a_{\beta-1} + b_{\beta-1} + d_{\beta-1},\\ &c_0 + d_0, c_1 + d_1, \cdots, c_{\beta-1} + d_{\beta-1},\\ &a_0 + c_0 + d_0, a_1 + c_1 + d_1, \cdots, a_{\beta-1} + c_{\beta-1} + d_{\beta-1},\\ &b_0 + c_0 + d_0, b_1 + c_1 + d_1, \cdots, b_{\beta-1} + c_{\beta-1} + d_{\beta-1},\\ &a_0 + b_0 + c_0 + d_0, \cdots, a_{\beta-1} + b_{\beta-1} + c_{\beta-1} + d_{\beta-1}),\end{aligned}$$

hence,

$$\begin{aligned}\sigma_2\Psi(w) = (&q_{\alpha-1} + r_{\alpha-1}, q_0, q_1, \cdots, q_{\alpha-1},\\ &q_0 + r_0, q_1 + r_1, \cdots, q_{\alpha-2} + r_{\alpha-2},\\ &a_{\beta-1} + d_{\beta-1}, d_0, d_1, \cdots, d_{\beta-1},\\ &a_0 + d_0, a_1 + d_1, \cdots, a_{\beta-2} + d_{\beta-2},\\ &a_{\beta-1} + b_{\beta-1} + d_{\beta-1}, b_0 + d_0, b_1 + d_1, \cdots,\\ &b_{\beta-1} + d_{\beta-1}, a_0 + b_0 + d_0, a_1 + b_1 + d_1, \cdots,\\ &a_{\beta-2} + b_{\beta-2} + d_{\beta-2}, a_{\beta-1} + c_{\beta-1} + d_{\beta-1}, c_0 + d_0,\\ &c_1 + d_1, \cdots, c_{\beta-1} + d_{\beta-1},\\ &a_0 + c_0 + d_0, a_1 + c_1 + d_1, \cdots, a_{\beta-2} + c_{\beta-2} + d_{\beta-2}\\ &a_{\beta-1} + b_{\beta-1} + c_{\beta-1} + d_{\beta-1}, b_0 + c_0 + d_0,\\ &b_1 + c_1 + d_1, \cdots, b_{\beta-1} + c_{\beta-1} + d_{\beta-1},\\ &a_0 + b_0 + c_0 + d_0, a_1 + b_1 + c_1 + d_1, \cdots,\\ &a_{\beta-2} + b_{\beta-2} + c_{\beta-2} + d_{\beta-2}),\end{aligned}$$

On the other hand,

$$\begin{aligned}\tau(w) = ((1+u)&x_{\alpha-1}, x_0, \cdots, x_{\alpha-2},\\ &(1+uv)y_{\beta-1}, y_0, \cdots, y_{\beta-2})\\ &= (r_{\alpha-1} + u(r_{\alpha-1} + q_{\alpha-1}), r_0 + uq_0, \cdots,\\ &r_{\alpha-2} + uq_{\alpha-2},\\ &a_{\beta-1} + ub_{\beta-1} + vc_{\beta-1} + uv(a_{\beta-1} + d_{\beta-1}),\\ &a_0 + ub_0 + vc_0 + uvd_0, \cdots,\\ &a_{\beta-2} + ub_{\beta-2} + vc_{\beta-2} + uvd_{\beta-2}).\end{aligned}$$

Hence,

$$\begin{aligned}\Psi\tau(w) = (&q_{\alpha-1}+r_{\alpha-1}, q_0, q_1, \cdots, q_{\alpha-2}, q_{\alpha-1},\\ &q_0+r_0, q_1+r_1, \cdots, q_{\alpha-2}+r_{\alpha-2},\\ &a_{\beta-1}+d_{\beta-1}, d_0, d_1, \cdots, d_{\beta-1}, a_0+d_0,\\ &a_1+d_1, \cdots, a_{\beta-2}+d_{\beta-2},\\ &a_{\beta-1}+b_{\beta-1}+d_{\beta-1}, b_0+d_0, b_1+d_1, \cdots, b_{\beta-1}+d_{\beta-1},\\ &a_0+b_0+d_0, a_1+b_1+d_1, \cdots, a_{\beta-2}+b_{\beta-2}+d_{\beta-2},\\ &a_{\beta-1}+c_{\beta-1}+d_{\beta-1}, c_0+d_0, c_1+d_1, \cdots, c_{\beta-1}+d_{\beta-1},\\ &a_0+c_0+d_0, a_1+c_1+d_1, \cdots, a_{\beta-2}+c_{\beta-2}+d_{\beta-2}\\ &a_{\beta-1}+b_{\beta-1}+c_{\beta-1}+d_{\beta-1}, b_0+c_0+d_0,\\ &b_1+c_1+d_1, \cdots, b_{\beta-1}+c_{\beta-1}+d_{\beta-1},\\ &a_0+b_0+c_0+d_0, a_1+b_1+c_1+d_1, \cdots,\\ &a_{\beta-2}+b_{\beta-2}+c_{\beta-2}+d_{\beta-2}).\end{aligned}$$

□

**Theorem 5.2.** *A $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive code $C$ is a $(1+u, 1+uv)$-additive constacyclic code if and only if $\Psi(C)$ is a binary quasi-cyclic code of index 2.*

*Proof.* If $C$ is $(1+u, 1+uv)$-additive constacyclic, then using Lemma(5.1) we have

$$\sigma_2(\Psi(C)) = \Psi(\tau(C)) = \Psi(C).$$

Hence, $\Psi(C)$ is a binary quasi-cyclic code of index 2. Conversely, if $\Psi(C)$ is a binary quasi-cyclic code of index 2, then using Lemma (5.1) again we get

$$\Psi(\tau(C)) = \sigma_2(\Psi(C)) = \Psi(C).$$

□

**Corollary 5.3.** *The image of a $(1+u, 1+uv)$-additive constacyclic code over $R_1^\alpha \times R_2^\beta$ under the Gray map $\Psi$ is a distance invariant binary quasi-cyclic code of index 2.*

# 6 Conclusion

In this paper, we studied $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive codes some property, including generator and parity check matrices for the codes. We fund the Gray map $\Phi$ is a distance preserving map and weight preserving map as well. At the end of this paper, we introduce the structure of $\mathbb{Z}_2[u]\mathbb{Z}_2[u,v]$-additive cyclic codes and constacyclic codes.

# References


[1] I. Aydogu, T. Abualrub and I. Siap, On $\mathbb{Z}_2\mathbb{Z}_2[u]$-additive codes, Interantional Journal of Computer Mathematics, 2105. vol.92, $No.9, 1806 - 1814$.

[2] I. Aydogu and Siap, The structure of $\mathbb{Z}_2\mathbb{Z}_{2^s}$-additive codes: Bounds on the minimum distance, Appl. Math. Inform. Sci. 7(6)(2013), pp. $2271 - 2278$.

[3] I. Aydogu, and I. Siap, On $\mathbb{Z}_{p^r}\mathbb{Z}_{p^s}$-additive codes, Linear and Multilinear Algebra, 2014

[4] J. Borges, C. Fernandez-Cordoba, J. Pujol, J. Rifa and M. Villanuena, $\mathbb{Z}_2\mathbb{Z}_4$-linear codes: Generator matrices and duality, Designs Codes Cryptogr. 54(2)(2010, ) pp. $167 - 179$.

[5] A. E. Brouwer, H. O. Hamalainen, P. R. J. Ostergard and N. J. A. Sloane, Bounds on mixed binary/ternary codes, IEEE Trans. Inform. Theory 44(1998), pp. 140-161.

[6] A. A. de Andrade and R. Palazzo, Linear codes over finite rings, TEMA Tend. Mat. Apl. Comput. $6(2)(2005)207 - 217$.

[7] M. EI-Alrash and M. AI-Ashker, Linear codes over $\mathbb{F}_2 + u\mathbb{F}_2$, Journal of the Islamie University of Gaza, $11(2), 53 - 68, 2003$.

[8] Eugene Spiegel, Codes over $\mathbb{Z}_m$, Information and control $35(1977)48 - 51$.

[9] A. R. Hammons, P. V. Kumar, A. R. Calderbank, N. J. A. Sloane and P. Sole, The $\mathbb{Z}_4$-linearity of Kerdock, preparata, Goethals and related codes, IEEE trans. Inform. Theory 40 (1994) pp.301-319.

[10] Liqi Wang, A family of constacyclic codes over $F_2 + uF_2 + vF_2 + uvF_2$, Journal of System Science and Complexity, October 2012.